

# THE INTERNATIONAL JOURNAL OF SCIENCE & TECHNOLEDGE

## Effect of Sugar Mill Effluents on Physico-chemical Properties of Soil of Doiwala Dehradun District of Uttarakhand (India)

**Bilal Bashir Bhat**

Junior Research Fellow, Department of Environment and Remote Sensing, Srinagar, Jammu and Kashmir, India

**Riyaz Ahmad Mir**

Junior Research Fellow, Department of Environmental and Remote Sensing, Srinagar, Jammu and Kashmir, India

**Mudasir Ahmad Dada**

Research Associate, Department of Environment and Remote Sensing, Srinagar, Jammu and Kashmir, India

**Shahzada Akther**

Lecturer, Department of Environmental Science, Jammu and Kashmir, India

### **Abstract:**

India is agriculture based country and there is a great demand in water for irrigation. Use of industrial effluent for agricultural and discharge of industrial effluent in the land of irrigation have become common practice in India as a result of which these toxic metals get accumulated in soil. The present study was aimed to investigate the impact of sugar industry on soil quality of area around the sugar industry of Doiwala Dehradun. Soil sampling was done from January to May from 04 different locations around a sugar industry. The various physico-chemical parameters were determined, including pH, Electrical Conductivity, Organic carbon, Potassium, Phosphorus and Sulphur. Results indicate that sugar mill effluent affected various soil parameters such as Organic carbon, Electrical conductivity, Sulphur and Potassium. Addition of nutrients by effluent improves the soil properties but the addition of too much nutrient like sulphur and potassium content in soil become toxic, decline the soil property.

**Keywords:** Sugar industry, Soil properties; effluent, Doiwala.




Figure 1: Location Map of Dehradun District

## 1. Introduction

India is agriculture based country and there is an excessive demand of water for irrigation while large quantity of effluent is discharged into water source as untreated. Use of industrial effluent for agricultural have become common practice in India, as a result of which these toxic metals get accumulated into plant tissue from the polluted soil. Disposal of industrial waste has created dual problem that has degraded the soil fertility and contaminated the food-chain. Sugar mill play a major role in polluting the water bodies and land by discharging a huge amount of wastewater as effluent. Several chemicals are used during the sugar manufacturing process mainly for coagulation of impurities and refining of the end products. Large amount of effluent generated during the manufacture of sugar contains a high amount of pollution load, particularly suspended solids, organic matters, press-mud, bagasses and air pollution. Discharge of sugar industry effluent to the land of irrigation influences the physico-chemical properties of soil.

Present study is conducted to identify the possible impact of sugar mill effluents on soil of Doiwala by analysis of various physico – chemical parameters of soil.

### 1.1. Study Area

Dehradun, also known as the Doon Valley, is situated in the state of Uttaranchal. Lying at the foothills of the Shivalik ranges in India, the city is said to have been founded by Guru Dronacharya (as per the Hindu Epic of Mahabharata).

Doiwala is a fast developing town of the state of Uttarakhandi along Haridwar – Dehradun. It is a proposed magnet mega city. City is growing as a transition town and works as central market for the nearby villages. It is a small town area surrounded by many small villages as Bhaniyawala, Jolly Grant, Laltappar, Chiddarwala, Teliwala, Hansuwala etc.



Figure 2

### 1.2. Sampling Sites

S#	Sampling Sites	Site
1.	Railway track drain	01
2.	Sugar mill field 1	02
3.	Sugar mill field 2	03
4.	Chand mari Agricultural field	04

Table 1

- Research site-I:-Railway track drain: - This site was located about 100 meters from sugar mill near at railway track Doiwala. Soil sample of the drain, which is close to railway track was selected as control site. The sediment taken was highly turbid and muddy black in appearance. The drain starts from sugar mill and passes through the slum area
- Research Site-II:-Sugar Mill Field:-This site was located about 200 meters from site-1.The soil sample of the agricultural field was selected as site-II.The agricultural field was influenced by human settlements. The site were the Soil was collected had wheat crop at germinating stage.The soil taken from this site was clayey in texture.
- Research Site III:-Sugar Mill Field 2:- It was located about 5 km down from the site II. The soil sample was taken from the agricultural field which was along the road side. The crop sown in such agricultural field was sugarcane. The soil sample taken from this site was sandy in texture.
- Research Site- IV:-Chandmari Agricultural field:- It was located about 1 km from the site III, near Chandmari village. The crop sown in this agricultural field was sugarcane. This field was also surrounded by heavy settlements. The soil taken from the site was clayey in the texture.

## 2. Material and Methods

All the samples were collected for sewage water of sugar mill and the nearby agricultural fields from four different Sites, situated at Doiwala Dehradun, 5 - 6 km from clocktower of Dehradun city [Site-I]. It was being irrigated with sewage water and agricultural practices are done in the different plots, and samples for Canal water irrigated land were collected behind railway tract of Doiwala [Site-II]. This site was located about 200 meters from site-I. The soil sample of the agricultural field was selected as site-II. The agricultural field was influenced by human settlements. The site were the Soil was collected had wheat crop at germinating stage. The soil taken from this site was clayey in texture. Site II: - It was located about 5 km down from the site II. The soil sample was taken from the agricultural field which was along the roadside. The crop sown in such agricultural field was sugarcane. The soil sample taken from this site was sandy in texture Site- IV:-Chandmari Agricultural field: - It was located about 1 km from the site III, near Chandmari village. The crop sown in this agricultural field was sugarcane. This field was also surrounded by heavy settlements. The soil taken from the site was clayey in the texture. The soil samples were collected from 3 different plots of each Site with the help of auger. Approximately about 500 gm soil was collected from a depth of 0-15 cm from each Site. The soil samples were taken and placed in a clean polythene bag. Two samples were collected from each Site. Various physico-chemical parameters of each land area were analyzed by different techniques in the laboratory.

S. No.	Physical parameters	November	January	May	Min. value	Max. value	Mean value
1	Colour	Grey	Grey	Grey	Grey		
2	Texture	Slity – clay	Slity – clay	Slity – clay			
A	Sand (%)	20	15	20	15	20	18.33
B	Clay (%)	60	65	55	55	65	60
C	Slit (%)	20	20	25	20	25	21.66
3	Porosity	0.0166	0.231	0.152	0.152	0.231	0.183
4	Conductivity ( $\mu\text{mho}$ )	0.095	0.205	0.023	0.095	0.205	0.107
5	Total dissolved solids (TDS)	109.2	151.9	112.9	109.2	151.9	124.66
<b>Chemical parameters</b>							
1	pH	7.5	8.0	7.6	7.5	8.0	7.7
2	Total Alkanlity (TA)	175	138	125	125	175	146
3	Chlorie (cl)	33.011	29.011	33.033	30.692		
4	Calcim (Ca++)	64.88	54.41	48.096	48.096	64.88	55.795
5	Magnesium (mg++)	54.09	54.09	44.26	34.90	54.09	44.416
6	Sodium (Na+)	19.00	24.00	27.00	19.00	27.00	23.33
7	Potassium (K <sup>+</sup> )	15.00	12.00	9.00	9.00	15.00	12.00
8	Phosphate – phosphorous (PO <sub>4</sub> – p) ) mg/L	0.038	0.50	0.21	0.21	0.50	0.186
9	Nitrate – nitrogen (NO <sup>3</sup> – N)	1.68	1.95	1.95	0.68	1.95	1.81
10	Organic matter (%)	3.67	3.62	5.32	3.62	5.32	4.20

Table 2: physico – chemical characteristics of Agricultural soil at Site 1

## 3. Result and Discussion

S no.	Physical parameters	November	January	May – 2011	Min. value	Max. value	Mean value
1	Colour	Dark grey	Dark grey	Dark grey			
2	Texture	clayey	clayey	Clayey			
3	Sand (%)	15	20	15	15	20	16.66
4	Clay (%)	60	75	65	60	75	66.66
5	Slit (%)	25	15	20	15	25	20
6	Porosity (%)	0.121	0.161	0.122	0.121	0.161	0.135
7	Conductivity (µmho)	0.127	0.258	0.222	0.022	0.258	0.136
8	Total dissolved solids (TDS)	154.8	120.1	157.2	120.1	157.2	144.33
<b>Chemical parameters</b>							
1	pH	8.5	7.2	7.5	7.2	8.5	7.733
2	Total Alkalinity (TA)	150	126	130	126	150	135.33
3	Chloride (cl <sup>-</sup> )	30.33	34.24	44.04	30.33	44.04	36.104
4	Calcium (Ca <sup>++</sup> )	72.4	68.41	32.06	32.06	72.4	57.623
5	Magnesium	47.6	28.26	29.2	28.23	47.6	35.02
6	Sodium (Na <sup>+</sup> )	30.00	27.00	30.00	27.00	30.00	29.00
7	Potassium (K <sup>+</sup> )	21.00	16.00	9.00	9.00	21.00	15.33
8	Phosphate – phosphorous (PO <sub>4</sub> – p )	0.061	0.45	0.012	0.012	0.061	0.039
09	Nitrate – nitrogen (NO <sup>3</sup> – N)	1.120	1.103	1.108	1.103	1.120	0.777
10	Organic matter (%)	4.99	4.99	7.06	4.99	7.06	5.68

Table 3: physico – chemical characteristics of Agricultural soil at Site – II

S#	Physical parameters	November	January	May – 2011	Min. value	Max. value	Mean value
1	Colour	Grey	Grey	Grey			
2	Texture	Sandy	Sandy	Sandy			
3	Sand (%)	70	75	75	70	75	73.333
4	Clay (%)	10	10	15	10	15	11.666
5	Slit (%)	20	15	10	10	20	15
6	Porosity (%)	0.102	0.141	0.112	0.102	0.141	0.118
7	Conductivity (umho)	0.758	0.165	0.013	0.013	0.758	0.312
8	Total dissolved solids (TDS)	98.8	113.7	69.9	69.9	113.7	94.133
<b>Chemical parameters</b>							
1	pH	8.8	7.6	7.8	7.6	8.8	8.066
2	Total Alkalinity (TA)	180	142	140	140	180	154
3	Chloride (cl <sup>-</sup> )	28.02	32.04	28.03	28.02	32.04	29.363
4	Calcium (Ca <sup>++</sup> )	42.0	51.42	26.03	26.03	51.42	39.816
5	Magnesium (mg <sup>++</sup> )	18.00	41.24	23.96	18.00	41.24	27.733
6	Sodium (Na <sup>+</sup> )	24.00	27.00	25.00	24.00	27.00	25.333
7	Potassium (K <sup>+</sup> )	14.00	19.00	9.00	9.00	19.00	14.00
8	Phosphate – phosphorous(PO <sub>4</sub> -P)	0.03	0.041	0.016	0.03	0.041	0.029
9	Nitrate – nitrogen (NO <sup>3</sup> – N)	1.129	1.060	1.075	1.060	1.129	1.088
10	Organic matter (%)	5.44	4.31	8.01	4.31	8.01	5.92

Table 4: physico – chemical characteristics of Agricultural soil at Site III.

S#	Physical parameters	November	January	May – 2011	Min. value	Max. value	Mean value
1	Colour	Grey	Grey	Grey			
2	Texture	Sandy	Sandy	Sandy			
3	Sand (%)	75.00	80.00	80.00	75.00	80.00	78.33
4	Clay (%)	10.00	10.00	10.00	10.00	10.00	10.00
5	Slit (%)	15.00	10.00	10.00	10.00	15.00	11.66
6	Porosity (%)	0.102	0.182	0.108	0.102	0.182	0.131
7	Conductivity ( $\mu\text{mho}$ )	0.839	0.068	0.016	0.016	0.839	0.308
8	Total dissolved solids (TDS)	200	153.2	101.6	101.6	200	151.6
<b>Chemical parameters</b>							
1	pH	8.2	7.8	7.8	7.8	8.2	7.933
2	Total Alkalinity (TA)	135	145	126	126	145	135.33
3	Chloride ( $\text{Cl}^-$ )	35.41	41.04	21.52	21.52	41.04	32.534
4	Calcium ( $\text{Ca}^{++}$ )	62.2	63.28	34.01	34.01	63.28	53.163
5	Magnesium ( $\text{Mg}^{++}$ )	17.8	19.41	24.99	17.8	24.99	20.733
6	Sodium ( $\text{Na}^+$ )	25.00	21.00	23.00	21.00	23.00	23.00
7	Potassium ( $\text{K}^+$ )	17.00	14.00	8.00	8.00	17.00	13.00
8	Phosphate – phosphorous ( $\text{PO}_4 - \text{p}$ )	0.041	0.083	0.011	0.011	0.083	0.045
9	Nitrate – nitrogen ( $\text{NO}_3 - \text{N}$ )	1.105	1.032	1.092	1.032	1.105	1.076
10	Organic matter (%)	4.76	3.67	6.92	3.67	6.92	5.11

Table 5: physico – chemical characteristics of Agricultural soil at Site IV

#### 4. Discussion

Physical chemical properties of soil samples was studied, all the samples are gray and dark grey in color and have unpleasant smell. The pH of soil is one of the most important physicochemical parameter. It affects mineral nutrient soil quality and much microorganism activity.

The present study is the first attempt to document the physico-chemical characteristics of agricultural soil of surrounding area of Doiwala sugar mill. The results of the investigation are detailed in the table.

The six months data on physico-chemical analysis of agricultural soil of surrounding area of Doiwala sugar mill at four sampling sites (site I – site IV) are presented in the table given later.

##### 1: Colour

Soil colour is important as it affects absorption and reflection of solar radiation and indicates chemical composition of soil. The colour of soil found in the catchment area of Doiwala sugar mill are grey. It was observed that the colour of soil found in open water zone was light grey in colour were as it was dark grey in vegetation covered zone.

##### 2: Texture

Texture is one of the most important physical properties of soils as it affects water retention nutrient availability, poor space, slope stability aeration and erosion susceptibility. In general soil was silty clay in texture were as sediment was sandy in texture. The composition of clayey was observed (10% -75%) in all the samples were as proportion of silt ranged from 5 % to 25 % and sand 20 - 80% respectively. Texture of soil influence the quality of soil and an excess of particle size may adversely affect the physical and chemical properties of soil required for vegetation growth (Donazzola et al 1984, Singh, 1997).

##### 3: pH

PH is one of the important parameter in water quality monitoring as it governs most of chemical reactions in water. At extremely high or low Ph values, many plants, animals and other aquatic biota may die and may pose hazardous impact on human and animal health.

The pH of the agriculture soil of the catchment area of Doiwala sugar mill in the present study was found to, be slightly alkaline to moderate throughout the study period. It was ranged from 7.5 -8.0 mg/100 g, 7.2 – 8.5 mg/100g, 7.6-8.8mg/100g and 7.8-8.2mg/100g in agricultural soil respectively from site I-IV. The average mean value of pH in agricultural soil ranged between 7.7 – 8.66 respectively.

##### 4.1. Conductivity

Conductivity is a measure of salt content of a sediment suspension and is therefore an indirect measure of osmotic pressure that plant roots must overcome for plant to take up water and nutrients. The conductivity of agriculture soil in the catchment area of Doiwala sugar mill varied from 0.095-0.205 mg /100g, 0.022-0.258 mg/100g, 0.013-, 0.758mg/100g and 0.016-0.839mg/100g in agricultural soil from site I to site IV respectively. The average mean value of conductivity in agricultural soil was ranged between 0.107-0.312.

Relatively higher conductivity observed in open water area of lake might be attributed to salt bearing materials probably reached to the lake sediment.

#### 4.2. Total Alkanity (T.A)

Total alkanity concentration was ranged between 125-175mg/100g in agricultural soil from site I to site IV respectively. The average mean value of alkanity in agricultural soil was 135.33-154.

Chloride:-

In the present study the chloride content in the soil and river sediments varied from 29.011-33.033mg/100g,30.033-44.04mg/100g,28.02-32.04 mg/100g and 21.52-41.4mg/100g in agricultural soil from site I to IV respectively. The average mean value of chloride in agricultural soil was ranged between 29.363-36.104.

#### 4.3. Porosity

The porosity of the agricultural soil sample of the catchment area of Doiwala sugar mill ranged from 0.152-0.231mg/100g, 0.121-0.161mg/100g,0.102 -0.141mg/100g and 0.102-0.182mg/100g respectively from I to IV. The average mean value of porosity in agriculture soil was ranged between 0.118-0.183.

Total dissolved solids (T.D.S):-

T.D.s in agricultural soil of the catchment area of Doiwala sugar mill ranged from 109.2-151.9mg/100g,69.9-113.7mg/100g and 101.6-200mg/100g respectively frpm site I to IV. The average mean value of T.D.S in agricultural soil was ranged between 94.133-151.6.

Calcium (ca):-

Calcium content in the soil ranged from 64.88mg/100g,32.06-72.4mg/100g,26.03-51.42mg/100g and 34.01-63.28mg/100g in agricultural soil from site I – IV respectively. The average mean value of calcium in agricultural soil was ranged between 39.816-57.623.

#### 4.4. Magnesium (Mg)

Magnesium content in the agricultural soil ranged between 34.90-54.09 mg/100g,28.26-47.6mg/100g,18.00-41.24mg/100g and 17.8 -24.99mg/100g in agricultural soil from site I \_ IV respectively. The average mean value of magnesium in agricultural soil was ranged between 20.733-44.416.

#### 4.5. Sodium (Na):-

The sodium content in soil was varied from 19.00-27.00 mg/100g,27.00-30.00mg/100g,24.00-27.00mg/100g and 21.00-23.00mg/100g in agricultural soil from site I to Iv respectively. The average mean value of sodium in agricultural soil was ranged between 23.00-29.00.

#### 4.6. Potassium (K+)

The value of potassium in agricultural soil of catchment area of Doiwala sugar mill ranged from 9.00-15.00mg/100g,9.00-21.00mg/100g,9.00-19.00mg/100g and 8.00-17.00 in agricultural soil from site I-IV respectively. The average mean value of potassium in agricultural soil was ranged between 12.00-15.33.

#### 4.7. Nitrate (No3-)

Agricultural wastes are generally organic materials. It includes crop residues, animal manure, poultry manure, fertilizers and pesticides. Due to increasing use of commercial fertilizers, effluents containing phosphate is being discharged into water which is then used for irrigation purposes in agricultural fields. The value of nitrate concentration of soil ranged from 1.068-1.095mg/100g,1.103-1.120mg/100g,1.060-1.129mg/100g and 1.032-1.105mg/100g from site I –IV respectively. The average mean value of nitrate in agricultural soil was ranged between 0.77-1.088.

#### 4.8. Phosphate-phosphorus (Po4-p)

The value of phosphorus in the agricultural soil of catchment area of Doiwala sugar mill ranged from 0.021-0.050mg/100g,0.012-0.061mg/100g,0.03-0.041mg/100g and 0.011-0.083mg/100g from site I-site IV respectively. The average mean value of phosphate-phosphorus in agricultural soil was ranged between 0.029-0.186.

#### 4.9. Organic Matter

Organic matter in agricultural soil or in catchment area of Doiwala sugar mill ranges from 3.62 – 5.32 mg/100g, 4.99 – 7.06 mg/100g, 4.31- 8.01mg/100g and 3.67 – 6.92 mg/100g respectively from site 1 to site IV. The average mean value of organic matter in agricultural soil was ranged between 4.20 – 5.92.

### 5. References

- i. Alexander, KEF (1971). Analyses of filter cake from South African sugar mills. Proc S Afr Sug Technol Assoc 45: 149-151.
- ii. Carter, MR (1986). Microbial biomass and mineralisable nitrogen in solonchic soils : influence of gypsum and lime amendments. Soil Biol Biochem 18: 531-537.
- iii. Chandra, R., K. Kumar and J. Singh: Impact of anaerobically treated and untreated (raw) distillery effluent irrigation on soil microflora, growth, total chlorophyll and protein contents of Phaseolus aureus L. J. Environ. Biol., 25: 381-385 (2004).

- iv. Chang, C.W.: Enzymic degradation of starch in cotton leaves. *Phytochemistry*, 21: 1263–1269 (1982).
- v. 12. Dunn, G.: A model for starch breakdown in higher plants. *Phytochemistry*, 13: 1341-1346 (1974).
- vi. 13. Dutta, S.K. and C.I. Boissya : Effect of paper mill effluent on growth yield and N, P, K contents of transplanted rice (*Oryza sativa* L. var. mahsuri). *J. Environ. Biol.*, 20: 29-32 (1999)
- vii. Gautam, D.D. and S. Bishnoi: Effect of dairy effluent on wheat (*Triticum aestivum*). *J. Ecology*, 4: 111-115 (1992).
- viii. Gupta, S., A.K. Pandey, N. C. Sharma, P. Pandey and C.D. Sharma: Studies on the biological treatment of digested distillery spent wash (2002).
- ix. Hackett, GAR, Easton, CA and Duff, SJB (1999). Composting of pulp and paper mill fly ash with wastewater treatment sludge. *Bioresource Technol* 70: 217-224.
- x. Haynes, RJ and Beare, MH (1996). Aggregation and organic matter storage in meso-thermal, humid soils. pp 213-262. In: MR Carter and BA Stewart (Eds). *Structure and Organic Matter Storage in Agricultural Soils*. CRC Lewis, Boca Raton.
- xi. Hosetti, B.B.: Treatment of sugar industry effluents by ponds and lagoons. *Environ. Biol.*, 16: 143-149 (1995).
- xii. Kaushik, A., K. Bela and C.P. Kaushik: Sugar mill effluent effect on growth photosynthetic pigments and nutrient uptake in wheat seedling in aqueous vs. soil medium, *Water Air Soil Pollution.*, 87: 38-46 (1996).
- xiii. Kisku, G.C., S.C. Barman and S.K. Bhargava: Contamination of soil and plants potentially toxic elements irrigated with mixed industrial effluent and impact in the environment. *Water Air Soil Pollute.*, 120: 121 – 137 (2000).
- xiv. Lerner, BR and Utzinger, JD (1986). Wood ash as a soil liming material. *Hort Sci* 21: 76-78.
- xv. McGarry et al., (1996). Effect of long term cane production on soil properties of a gleyed podzolis soil near Bundaberg, Queensland.
- xvi. Metcalf and Eddy: Third edition. *Waste water engineering, treatment, disposal and Reuse*, Tata McGraw Hill Publication Limited, New Delhi. pp. 633-704 (1995).
- xvii. Misra, R.: Effect of Urbanization and industrialization on the natural vegetation of India. A.K. Sharma and A. Sharma (Eds.) *Impact of the development of science and technology on environment*. Indian Science Congress Association Publication, Calcutta, 5-38 (1981).
- xviii. Moberly, PK and Meyer, JH (1978). An evaluation of poultry manure as a sugarcane fertilizer. *Proc S Afr Sug Technol Assoc* 45: 136-141.
- xix. Nath, K. and Y.K. Sharma: Effect of sugar factory effluent on seed germination and seedling growth in pulses. *Boil. Memoirs*, 28: 73-78 (2002).
- xx. Nath, K., D. Singh and Y. K. Sharma: Assessment of seed performance under the influence of distillery effluent. *J. Ecophysiol. Occup. Hlth.*, 7: 55-62 (2007a).
- xxi. Om, H., N. Singh and M.S. Arya : Combined effect of wastes of distillery and sugar mill on seed germination, seedling growth and biomass of okra (*Abelmoschus esculentus* L. Moench). *J. Environ. Biol.*, 15: 171-175 (1994).
- xxii. Raison, RJ (1979). Modification of the soil environment by vegetation fires, with particular reference to nitrogen transformations: a review. *Plant Soil* 51: 73-108.
- xxiii. Ramana, S., A.K. Biswas, S. Kundu, J.K. Saha and R.B.R. Yadava : Efficacy of distillery effluent on seed germination and seedling growth in mustered cauliflower and radish. *Proc. Nat. Acad. Sci. India*, 71: 129-136 (2001).
- xxiv. Roth, G (1971). The effects of filter cake on soil fertility and yield of sugarcane. *Proc S Afr. Sug Technol Ass* 45: 142-148.
- xxv. Sharma, V., R. Sharma and K.D. Sharma : Distillery effluent effect on seed germination, early seedling growth and pigment content of sugarbeet (*Beta vulgaris* L. var. Mezzanau–Poly). *J. Environ. Biol.*, 23: 77-80 (2002).
- xxvi. Srivastava, P.K., G.C. Pandey and S. Naraliya : Bioaccumulation of distillery effluent metals by some aquatic macrophytes. *Proc. Nat. Acad. Sci. India*, 70: 311-318 (2000).
- xxvii. Swaminathan, K., J. Aryan, R. Gurusamy: Effect of glucose factory effluents on seed germination and seedling development of Groundnut (*Arachis hypogea* L.). In: R.C. Dalela (Eds) *Environ. Impact on Biosystems*. The Academy of Environ. Biol., 81-87(1989).
- xxviii. Thevenot, C., C. Lauriere, C. Mayer, Cote-Simond and J. Daussant : a amylase changes during development and germination of maize kernels. *J. Plant Physiol.*, 140: 61 – 65 (1992).
- xxix. Vance, ED (1996). Land application of wood-fired and combination boiler ashes: an overview. *J Environ Qual* 25: 937-944.
- xxx. Veer, B. and K. Lata: Effect of polluted municipal waste water on growth, yield and levels of some biochemical components of *Triticum aestivum* L. cv. SD 2204. *Acta Bot. Indica*, 15: 242-246 (1987).
- xxxi. Zalawadia, N.M., R.G. Patil and S. Raman : Effect of distillery wastewater with fertilizer on onion and soil properties. *Indian J.Soc.So. Sci.*, 44: 802-804 (1996).
- xxxii. Zhang, F (2001). Application of waste ashes to agricultural land; effect of incineration temperature on chemical characteristics. *SciTotal Env* 264: 205-214.