

THE INTERNATIONAL JOURNAL OF SCIENCE & TECHNOLEDGE

Profile of Hearing Impairment and its Effect on Life Style of Sample Population in the Medinipur Sadar Subdivision, Paschim Medinipore District, West Bengal, India

Kausik Chatterjee

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India
Molecular Medicine Laboratory, Department of Bio-Medical Laboratory Science & Management Vidyasagar University,
Midnapore, West Bengal, India
Department of Physiology, Midnapore College, Midnapore, West Bengal, India

Sanatan Hazra

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India

Sanjay Kumar Kundu

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India

Soumyajit Maiti

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India
Molecular Medicine Laboratory, Department of Bio-Medical Laboratory Science & Management Vidyasagar University,
Midnapore, West Bengal, India

Hanif Sikdar

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India

Souvik Goswami

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India

Gita Pal

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India

Debidas Ghosh

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India
Molecular Medicine Laboratory, Department of Bio-Medical Laboratory Science & Management Vidyasagar University,
Midnapore, West Bengal, India

Dipak Kumar Bhuniya

Audiology Division of Midnapore Rehabilitation Centre for Children
(Rehabilitation Council of India, Govt. of India Recognized Centre), Midnapore, Midnapore, West Bengal, India
Department of Physiology, Midnapore College, Midnapore, West Bengal, India

Abstract:

To investigate the prevalence of hearing impairment and its effect on life style of the sample population in Medinipur Sadar Subdivision, Paschim Medinipore District present cross-sectional study was designed. Current study recruited 700 respondents from this subdivision were used in the data collection. The procedure adopted included interviews, questionnaires, and physical examination at Audiology Division of Midnapore Rehabilitation Centre for Children (Rehabilitation Council of India, Govt. of India recognized centre), Midnapore. Out of the 700 respondents, 103 (representing 14.71%) were diagnosed with significant hearing impairments and majority of the respondents were men than women had mild hearing loss with occurrence of conductive hearing impairment more than other types of hearing impairment. Presbycusis and noise were the major causes of the sensorineural hearing impairment alongside ear wax, otitis media, and noises induce factors the other frequent causes. Hearing impairment was found with significant impact on normal life style. These findings have significant implications on the need of resource development for prevention and rehabilitation.

Keywords: *Audiology, conductive hearing loss, presbycusis, sensorineural hearing impairment, Midnapore sadar subdivision*

1. Introduction

Hearing impairment a growing world wide burden and became a principal public health task in the world (WHO, 1995);. The World Health Organization (WHO) reported an escalating trend in the statistics of global number of persons with hearing impairment is most likely due to a mixture of improved diagnosis, early detection, old age, ototoxic drugs and long-term exposure to environmental noise (Crofton et al., 1994). It has been showed that globally out of 278 million hearing impaired people 89% are adults while 11% were children under 15 years (WHO, 2014; Smith,2008), and using of hearing aids meet up less than 10% of global need. So, it has profound individual, societal, and economic consequence and in high-income countries, hearing impairment has been shown to have very large financial cost and these are likely in low and middle-income countries (Smith 2008). In developing countries a widespread insufficiency of appropriate data about hearing impairment still exists (UNICEF, 2013; Groce, 2004), which is urgently needed in order to raise awareness, for intervention, calculate about the needs, monitoring of outcome, economic analyses. On that background our aim was to investigate out the profile and demographic pattern of hearing impairment along with its causative historical background i.e. family history, past medical history and drugs history and present factors predispose respondents in the Medinipur Sadar subdivision of Paschim Medinipore district.

2. Materials and Methods

To establish the impact database required for future needs and developments seven hundred (700) people, with both of male and female subjects were incorporated in this study. The Paschim Medinipore district comprises four sub divisions: Kharagpur, Ghatal, Jhargram and Medinipur Sadar, which is consists of Midnapore municipality and six community development blocks: Medinipur Sadar, Garhbeta-I, Garhbeta- II, Garhbeta-III, Keshpur and Shalboni. The sampling design was cluster random sampling (Ahmed, 2009) along side cross-sectional research design (Lee, 1994) was employed for this study. A ratio of the total sample was selected from clusters by a random technique to obtain the required total sample. The research instruments employed were all regulated tools with testing its validity and reliability. The research tools consist of the following: i) Questionnaire: Pre-tested open and closed ended questions questionnaire was used to obtain respondents, opinions. The questionnaires enabled the researcher to collect data from a large number of respondents ii) Interview: To describe the uniqueness of the study participants possible causes of common ear infection, hearing impairment, demographic distribution and its profile were recorded through interview of the subject. iii) Physical Ear Examination: After the interview the participants were subjected to physical ear examination by using otoscopy to determine the presence or absence of outer, middle and inner ear infections. Participants with any form of ear infections were not be permitted to the audiometric examination iv) Audiometry: The ELKON [Model- eda 3 N 3 mille] audiometer was used calibrated to the ANSI standard (ASHA, 1978). The purpose of the screening was first explained to them and date for the screening was arranged in advance. A separate room with an average noise level of 41dB SPL was selected as test room (Frank and William, 1993). On their day of screening, subjects were brought into the test rooms in group of twenty and instructed regarding the procedure for the hearing test, asked about any ear discharge or earache and was presented with questionnaire. Audiometric screening was carried out at three frequencies 1000Hz, 2000Hz and 4000Hz (OSHA,1983). Ambient noise level in the test room limited testing to these mid-range and high frequencies. The degree of hearing impairment was based on the criteria developed by the world health organization i.e. for each ear, "pass" was operationally defined as responding properly to stimuli at 30dB HTL and at all three frequencies. When a respondent failed to respond at any of these frequencies, the tone was re-presented at 35, 40 and 45 dB HTL followed by the recording of response (Martin,1986).

3. Results and Discussion

The Table 1 showed that among the 700 respondents selected from the Medinipur Sadar Subdivision, 424 of the respondents (60.57%) came from Medinipur Sadar. 46 (6.57%), 38 (5.43%) and 51 (7.29%) of the respondents came from Garhbeta-I, Garhbeta-II and Garhbeta-III respectively. 106 of the respondent (15.14%) came from Shalboni community development blocks.

Community development blocks	Frequency	Percentage (%)
Medinipur Sadar	424	60.57
Garhbeta-I	46	6.57
Garhbeta-II	38	5.43
Garhbeta-III	51	7.29
Keshpur	35	5.00
Shalboni	106	15.14
TOTAL	700	100

Table 1: Geographical Area of Respondents within the Medinipur Sadar Subdivision

Among 700 respondents in the target subdivision, 239 (39.83%) were male while 361(60.17 %) were females (Table 2).

Gender	Frequency	Percentage
Male	280	40
Female	420	60
TOTAL	700	100

Table 2: Distribution of gender among respondents

The age distributions of the respondents in the present study are shown in Figure 1. Out of 700 respondents, 175 (25 %) belonged to the 0- 20 age group, 126 (18%) were 21-40 years old, 238 (34%) under the 41-60 age group and 161 (23%) were above 60 years.




Figure 1: A bar chart illustrating the age distribution of the respondents in the study area

Out of 700 respondents selected from the said area, majority of the respondents (46.43%) were working in the non-formal sector with 15.71% of them in the formal sector and 37.86% of the respondents were unemployed (Table 3).

Employment status		Number and Percentage (%) (n=700)	
Employed	Formal profession (Govt. sectors like education, finance, health, engineering, environmental and agricultural sectors)	110 (15.71%)	435(62.14%)
	Non-formal profession (Mechanic, cotton mill-worker, chain-shaw mill worker, car driver, constructional worker)	325 (46.43%)	
Un-employed		265 (37.86%)	

Table 3: Profile of employment status of respondents in Medinipur Sadar Subdivision

Table 4 showed the age distribution of the results of the field audiometric screening test. To fail field audiometric screening test means not able to respond correctly to stimuli at 30dB HTL for all frequencies and to pass audiometric screening test means to respond correctly to stimuli at 30dB HTL for all frequencies. Out of the 700 respondents selected, majority of the respondents 515 (73.57%) passed the audiometric screening test and were not referred. Meanwhile, 185 (26.43%) of the respondents failed the audiometric screening test and were accordingly referred to MRCC for further clinical examination and management. Out of 175 respondents belonged to the 0-20 age 46 respondents failed the audiometric screening test while 129 passed the audiometric screening test. From 126 respondents under the 21–40 age groups, 17 respondents failed the audiometric screening test as 109 respondents passed the audiometric screening test. Of the 238 respondents belonged to the 41–60 age group, 57 respondents failed the audiometric screening test even as 181 respondents passed the audiometric screening test. Out of 161 respondents above 60 years, 65 respondents failed the audiometric screening test whereas 96 respondents passed the audiometric screening test.

Age (Years)	Passed screen test	Failed in screen test	Total
Below 20	129 (73.71%)	46 (26.29%)	175
21-40	109 (86.51%)	17 (13.49%)	126
41-60	181 (76.05%)	57 (23.95%)	238
Above 60	96 (59.63%)	65 (40.37%)	161
Total	515 (73.57%)	185 (26.43%)	700

Table 4: Results of audiometric screening

Table 5 shows the results of pure tone audiometry and threshold levels (Martin,1986) for the current study. Out of 185 respondents who failed screening and were referred for further evaluation 103 respondents had hearing impairment at the low (250 and / or 500 Hz) and high frequencies (4000 and /or 8000Hz). Twenty (29) respondents with hearing impaired, having a threshold range of 26dB-61 dB, 9 respondents hearing impaired, having threshold range of 26 dB - 40 dB, 21 respondents were hearing impaired, having a threshold range of 26 dB - 42 dB and 44 respondents were hearing impaired, having a threshold range of 26 dB-85 dB.

Threshold level of Hearing impaired (dB)	Failed in screen test and referred	Detected as hearing impaired
26-61	46	29
26-40	17	9
26-42	57	21
26-85	65	44
Total	185	103

Table 5: Results of pure tone audiometric and threshold levels

Table 6 showed the pattern of hearing impairment among the respondents. Out of 103 respondents with hearing impairment, 46 (44.66%) had problems in the right ear, 38 (36.89%) had problems in the left ear and 19 (18.45%) had problems in both ears.

Hearing impairment pattern	Number and percentage (%) (n=103)
Right ear	46 (44.66%)
Left ear	38 (36.89%)
Both ear	19 (18.45%)

Table 6: The pattern of hearing impairments

The types of hearing loss of respondents using tuning fork in the recent study were shown in Table 7. Out of 103 respondents with hearing loss, 97 (72%) had conductive hearing loss, 32 (24%) had sensorineural hearing loss and 6 (4%) had mixed hearing loss.

Types of Hearing impaired	Number and percentage (%) (n=103)
Conductive hearing loss	68 (66.02%)
Sensorineural hearing loss	31 (30.10%)
Mixed hearing loss	4 (3.88%)

Table 7: Types of hearing impairment

The geographical area of respondents with hearing impairment in the present study is expressed in Table 8. Out of 103 respondents with hearing impairment, 41 of respondents (39.80%) came from Medinipur sadar; 12,6,14 of the respondents (11.65,5.83,13.59%) came from Garhbeta-I, II, III; 10 of the respondents (9.71%) came from Keshpore and 20 of the respondents (19.42%) came from Shalboni community development blocks. The higher number of hearing impairment in Midnapore sadar in comparison to others may due to the number of population in each block. It was however observed that the higher the population, there were more hearing problems (Olusanya and Okolo, 2006). The higher percentage of hearing impairment could be attributable to the load of noise in Midnapore sadar.

Community development blocks	Frequency and percentage (%)
Medinipur Sadar	41(39.80%)
Garhbeta-I	12(11.65%)
Garhbeta-II	6(5.83%)
Garhbeta-III	14(13.59%)
Keshpur	10(9.71%)
Shalboni	20(19.42%)
TOTAL	103(100%)

Table 8: Geographical area of respondents with hearing impairments

The age distributions of the respondents with hearing impairment in the current study were shown in Table 9. Out of 103 respondents, 28 (27.18 %) belonged to the 0-20 age group, 11 (10.68 %) were 21- 40 years old, 19 (18.45 %) under the 41-60 age group and 45 (43.69%) were above 60 years. The results mean that respondents above 60 years had higher hearing impairment, resembles the findings of Robinson et al. (1979) that one in five adults over the age of 80 suffer from age related hearing loss.

Age (Years)	Number and percentage
Below 20	28 (27.18%)
21-40	11 (10.68%)
41-60	19 (18.45%)
Above 60	45 (43.69%)
Total	103 (100%)

Table 9: The age distribution of respondents with hearing impairment

The gender distribution of respondents with hearing impairments in the present study are shown in figure 2. Out of 103 respondents with hearing impairments, 42 (59%) were female while 61 (41%) were male. So, it could be said that males in Medinipur Sadar subdivision are more exposed to high risk employment such as jobs where noisy equipments are used (Miller, 1971).




Figure 2: A pie chart illustrating the gender distribution of respondents with hearing impairments

The medical conditions attributed to hearing impairment in respondents are shown in Table 10. Out of 68 respondents with conductive hearing impairment, 33(48.53%) had wax, 10 (14.71%) had otitis media, 23 (33.82 %) had presbycusis and 2 (2.94 %) were taken ototoxic drugs (Roeser et al.,2005; Franks and Morata,1996). Olusanya et al (2004) reported that among school children screened with otoscopy had unilateral or bilateral impacted wax occluding the tympanic membrane leads to hearing loss also supported by our study. Presbycusis recorded the next highest medical condition common in said subdivision. It tends to agree with the findings of Robinson and Sutton (1979) that one in five adult over the age of 80 suffer from age-related hearing loss.

Conditions	Number and percentage (n=68)
Wax	33 (48.53%)
Otitis media	10 (14.71%)
Presbycusis	23 (33.82%)
Drug toxicity	2 (2.94%)

Table 10: Medical conditions attributable to hearing impairment in respondents

From the 103 respondents, 21 (20.39%) and 6 (5.83 %) had family history of hearing impairment and speech disorders. Family history of mental retardation or cerebral palsy and learning disability was carried by 2 (1.94%) and 5(4.85%) of the respondents. Three (2.91%) had family history of visual impairments and 66 (64.08 %) had none of these family histories (Table 11). This is in line with the findings of Schraders et al. (2010) and White (2004) that if a family had a dominant gene for deafness it would persist across generations because it would visible itself in the offspring even if it is inherited from only one parent. Schraders (2010) also claimed that if a family had genetic hearing impairment caused by a recessive gene it will not always be apparent as it will have to be passed onto offspring from both parents.

Family history	Number and percentage
Hearing impairment	21 (20.39%)
Speech disorders	6 (5.83%)
Mental retardation or cerebral palsy	2 (1.94%)
Learning disability	5 (4.85%)
Visual impairment	3(2.91%)
None	66(64.08%)
Total	103 (100%)

Table 11: Profile of family history of the hearing impaired subjects

The medical histories of the respondents with hearing impairment showed in Table 12. Among 103 respondents with hearing impairment 9(8.74%) suffered from childhood measles, 2 (1.94 %) had cerebrospinal meningitis, 5 (4.85 %) had diabetes mellitus, 13 (12.62 %) had medical history of hypertension and 74 (71.85%) had no such type of previous medical history. The findings are in consistence with literature of Shargorodsky et al, (2010) that medical history by itself is not associated with an overall greater risk of hearing loss.

Medical history	Number and percentage
Childhood measles	9 (8.74%)
Meningitis	2 (1.94%)
Diabetes mellitus	5 (4.85%)
Hypertension	13 (12.62%)
None	74 (71.85%)
Total	103 (100%)

Table 12: Profile of medical history of the respondents

Table 13 showed the profile of medication history of respondents with hearing impairment. Out of 103 respondents, 9 (8.74%) are taking or had taken antibiotics, 2 (1.94%) and 5 (4.84%) had taken antidiabetics and antihypertensive drugs, other types of drugs was taken by 13 (12.62 %) respondents and 74 (71.85 %) had none of these drug histories. Our finding tends to agree with the claim of Fukushima (2004); Matz and Naunton, (1968); Robinson and Cambon (1964) who said that the drugs are known to common cause hearing impairment.

Medication history	Number and percentage
Antibiotics drug	9 (8.74%)
Antidiabetic drug	2 (1.94%)
Antihypertensive drug	5 (4.85%)
Others drugs	13 (12.62%)
None	74 (71.85%)
Total	103 (100%)

Table 13: Profile of medication

The occupational conditions attributed to hearing impairment in respondents are shown in Table 14. Out of 103 respondents with occupational conditions of hearing impairment, 23 (22.33%) were formal profession, 75 (72.82%) respondents were attached with non-formal profession and 5 (4.85 %) personnel were unemployed. This finding correlates with the claim of WHO that 16% of deafness is due to occupational noise (WHO, 2004). It has been observed that mill operators in India exposed to noise levels exceeding 90dB. Hence mill operators have evident of hearing loss (Sataloff and Sataloff, 1993). Alongside males exposed to noise in farm jobs where noisy equipment such as tractors, chainsaws, and grain crushers are used also suffers from the same.

Employment status		Number and Percentage (%) (n=103)	
Employed	Formal profession (Govt. sectors like education, finance, health, engineering, environmental and agricultural sectors)	23 (22.33%)	98(95.15%)
	Non-formal profession (Mechanic, cotton mill-worker, chain-shaw mill worker, car driver, constructional worker)	75 (72.82%)	
Un-employed		5(4.85%)	

Table 14: Hearing impairment associated with occupational status

Among 103 people with hearing impairment, 22 (21.36%) had no formal education, 41 (39.81%) had primary education, 31(30.10%) had up to secondary education while 31 (30.10%) of them attained up to higher secondary and 9 (8.74%) of the respondents had education above higher secondary. This result shows that most people suffering did not achieve higher education. It may be said that they had been marginalized by the society and caused many of them to stop attending school. Another likely factor leads to drop out of school may also be due to insufficient sign language professionals in the schools in that subdivision (Carney and Moeller, 1997), which has significant impact on social development (Table 15).

Educational status	Number and percentage
No formal education	22 (21.36%)
Up to primary education	41 (39.81%)
Up to secondary education	31 (30.10%)
Above higher secondary education	9 (8.74%)
Total	103 (100%)

Table 15: Educational Status of Respondents with Hearing Impairments

The findings showed that out of the 103 respondents with hearing impairments, a major part 33 (32.04%) were separated or divorced whereas 19 (18.45%) of them were single. Thirty three (32.04 %) of the respondents was married. Therefore, once majority of them were divorced or separated, it is likely that their hearing impairment has influenced their married. This supposedly should mean their inability to communicate to each other and this has destroyed their relationship especially those who do not know the sign language resembles findings of other also (Hétu et al.1993) (Table 16)

Marital Status	Number and Percentage (%) (n=103)
Un-married	19 (18.45%)
Married	51 (49.52%)
In separation status or divorce	33 (32.04%)
Total	103 (100%)

Table 16: Marital Status of respondents with hearing impairments

4. Conclusion

The purpose of this study was to survey the prevalence of hearing impairment in the Medinipur Sadar Subdivision of Paschim Medinipore District, West Bengal. The data has shown that out of 700 respondents screened for hearing impairments, 103 were diagnosed as having significant hearing impairment, which has a significant impact on the district. Again, the results showed that the frequent causes of hearing impairment were as follows: wax, presbycusis, otitis media and noise induce factors as in the case of machine operators in mill and factory operators. It was moreover found out that medicine also caused hearing impairment. Besides it was also revealed that families with dominant genes for hearing impairment have the tendency of spreading the disease across generations. Hearing impairment had detrimental effect on education acquirement process and normal married life. It can be concluded that most of these findings is preventable if appropriate measures is taken.

5. References

1. Crofton, K.M., Lassiter, T. L. & Robert, C. S. (1994). Solvent Induced Ototoxicity in Rats: An Atypical Selective Mid-Frequency Hearing Deficit. *Hearing Research*, 80, 25-30.
2. WHO (World Health Organization). (1995). Prevention of Hearing Impairment. Resolution of the 48th World Health Assembly, WHA48.9. Geneva.
3. Smith, A. (2008). Demographics of Hearing Loss in Developing Countries. *Audiology in Developing Countries*. Nova Science Publisher, Inc., New York, USA. Pp.21-47.
4. WHO (World Health Organization). (2014). Deafness and Hearing Loss. Fact Sheet No. 300. Geneva.
5. UNICEF. (2013). Children and Young People with Disabilities. Fact Sheet. UNICEF/BANA 2007-00655.NY,USA.1-36
6. Groce, N.E. (2004). Adolescents and Youth with Disabilities: Issues and Challenges. *Asia Pacific Disability Rehabilitation Journal*. 15:13-32.
7. Ahmed, S. (2009). Methods in Sample Surveys.140.640.Cluster Sampling. Dept. of Biostatistics School of Hygiene and Public Health Johns Hopkins University.Pp.1-14.
8. Lee, J. (1994). Odds Ratio or Relative Risk for Cross-Sectional Data?. *International Journal of Epidemiology* 23: 201–203.
9. American Speech-Language- Hearing Association. (1978). Guidelines for manual pure-tone threshold audiometry. *ASHA*, 20, 297–301.
10. Frank, T. & Williams, D.L. (1993). Ambient noise levels in audiometric test rooms used for clinical audiometry. *Ear Hear*; 14:414-422.
11. Occupational Safety and Health Administration. (1983). Occupational Noise Exposure: Hearing Conservation Amendment. *Federal Register*. 48, 9738-9783.
12. Martin, FN.(1986). Introduction to Audiology, 5th edition. Prentice-Hall Inc., Englewood Cliffs, New Jersey.
13. Robinson, D.W. & Sutton G.J. (1979). Age Effect in Hearing-A Comparative Analysis of Published Threshold Data. *Audiology*. 1979; 18: 320-334.
14. Olusanya, B. & Okolo, A.A. (2006). Early Hearing Detection at Immunization Clinics in Developing Countries. *International Journal of Pediatric Otorhinolaryngology*. 70: 1495-1498.
15. Miller, J.D. (1971) Effects of Noise on People. U.S. *Environmental Protection Agency* Publication No. NTID 300.7 Washington, D.C, Pp.93.
16. Carney, A. E., & Moeller, M. P. (1997). Treatment Efficacy: Hearing Loss in Children. *Journal of Speech, Language, and Hearing Research*. 41: S61-S84.
17. WHO (World Health Organization). (2004). Guidelines for Hearing Aids and Services for Developing Countries.2nd Ed (WV 274). Geneva. Pp.1-26
18. Roeser, R.J., Lai, L. & Clark, J.L. (2005) Effect of Ear Canal Occlusion on Pure Tone Threshold Sensitivity. *Journal of the American Academy of Audiology*. 16:740- 746.
19. Franks, J.R. & Morata, T.C. (1996). Ototoxic Effects of Chemical alone or in Concert with Noise: A Review of Human Studies. In: Axelsson, A., Borchgrevink, Hamernik RP, Hellström PA, Henderson D, Salvi R (Eds.), *Scientific Basis of Noise-Induced Hearing Loss*, New York: Thieme. pp.437-472.
20. Sataloff, R.T. & Sataloff, J. (1993). *Occupational Hearing Loss*, 2nd edition, Marcel Dekker, Inc., New York, NY.
21. Olusanya, B.O., Okolo, A. A. & Adeosun, A. A. (2004). Predictors of hearing loss in school entrants in a developing country. *Journal of Postgraduate Medicine*.50:173-179.
22. Schraders, M., Oostrik, J., Huygen, P.L., Strom, T.M., vanWijk, E., Kunst, H.P., Hoefsloot, L.H., Cremers, C.W., Admiraal, R.J. & Kremer, H. (2010). Mutations in PTPRQ are a cause of autosomal-recessive nonsyndromic

- hearing impairment DFNB84 and associated with vestibular dysfunction. *The American Journal of Human Genetics*. 86:604–610.
23. Shargorodsky, J., Curhan, S.G., Roland, E. & Curhan, G. C. (2010) A prospective study of cardiovascular risk factors and incident of hearing loss in men. *Laryngoscope*. 120:1887-1891.
 24. White, K.R. (2004). Early hearing detection and intervention programs: Opportunities for genetic services. *America Journal of Medical Genetics A*, 130A, 29- 36.
 25. Fukushima, H. (2004). The effects of type I diabetes mellitus on the cochlear structure and vasculature in human temporal bones. *The National Temporal Bone Registry*, Summer 2004 issue, Volume 12, #1.
 26. Robinson, G.C. & Cambon, K.G. (1964) Hearing loss in infants of tuberculous mothers treated with streptomycin during Pregnancy. *The New England Journal of Medicine*. 271:949-951.
 27. Héту, R., Jones, L. & Getty L.(1993).The Impact of Acquired Hearing Impairment on Intimate Relationships: Implications for Rehabilitation. *International Journal of Audiology*. 32: 363-380